

Researches concerning iron bioavailability in complex foods

Corina CIOBANU and Natalia MOTRUC

Technical University of Moldova, 168 Stefan cel Mare St., MD-2004 Chisinau, Republic of Moldova
Phone: +373 22 237861, Fax: +373 22 235441, E-mail: corinaciobanu@mail.ru, motruc_n@mail.ru

Abstract

Iron bioavailability in fortified products was researched *in vivo* conditions. It is established, that the consumption of iron fortified bread products promotes improvement of the basic haematological indices. In the presence of induced experimental anaemia, the serum iron concentration was restored entirely in groups that received iron fortified sourdough bread, and in proportion of 45-45 % in the case of groups that received iron fortified bread prepared by the diphase procedure. On the basis of the received experimental results it has been elaborated an algorithm, allowing to determine the iron absorption coefficient from a representative meal and the estimation of the providing degree of nutritional requirements for different groups of population.

Key words: fortified bread, haematological indices, experimental anaemia, serum iron concentration, iron absorption coefficient, acid-lactic fermentation.

Rezumat

Biodisponibilitatea fierului din produse fortificate a fost cercetată *in vivo*. În condițiile unui statut normal în fier n-a fost atestată nici o influență a metodei de panificație asupra indicilor hematologici ai grupelor examinate, factorul principal fiind conținutul de fier din regim. În prezența anemiei experimentale la grupa care a primit în alimentație pâine fabricată prin metoda cu o etapă de fermentație acido-lactică, concentrația fierului seric s-a restabilit complet pe durata experimentului, iar la grupa alimentată cu pâine, preparată prin metoda tradițională (bifazică) concentrația fierului seric s-a restabilit doar în proporție de 40-45%. Pe baza rezultatelor experimentale a fost elaborat un algoritm, care permite determinarea coeficientului de absorbție al fierului dintr-o masă reprezentativă și estimarea gradului de acoperire a cerințelor nutriționale pentru diferite grupuri ale populației.

Cuvinte cheie: pâine fortifiată, indici hematologici, anemie experimentală, concentrația fierului seric, coeficient de absorbție a fierului, fermentație acido-lactică.

1. Introduction

According to WHO criteria, anaemia in the Republic of Moldova represents a major public health problem (UNICEF 1996–2000). The results of the investigations drawn under UNICEF supervision over a group of 792 children under 5 years old and their mothers show anaemia presence at 47% of children between 6 and 12 months old, at 28% of children under 5 years old and at 40% of women of fertile age (UNICEF 2002). According to the statistics of the Ministry of Health, anaemia rate in the case of children till one year old constitutes 20%. It is common knowledge that one anaemia case corresponds to one case of iron nutritional deficiency, it is possible to assume that

approximately half of the children under 5 years old and 40% of fertile age women have iron deficiency (Motruc *et al.*, 2005). Prevalence of blood and haematopoietic system diseases increased during last 5 years with 46.4% and for anaemia with 50.3%.

One of largely used methods in the highly developed countries is food fortification with deficient micronutrients (Bauernfeind, 1988). This method is easily realized and has very low cost in the case of the mineral micronutrients (Luten *et al.*, 1996). Refining of cereal products leads to an important loss of minerals and fibres and may have important consequences on such disease as colon cancer, diabetes mellitus and heart diseases. Flour fortification with micronutrients represents an

outspreed procedure which is aiming the reinstatement of micronutrient composition affected during processing. Thus, if non milled wheat contains approximately 33 mg/kg iron, after milling this amount is reduced to 11 mg/kg. Administration of 20 mg/kg iron to the flour will confer its natural qualities.

However, food fortification cannot be considered as a simple mechanical administration of the supplements (Turk *et al.*, 1996). In the case of vegetal products or combined nutrition (animal origin products consumed together with the cereals) the main cause for the demineralization is considered to be phytic acid (myo-inositol hexafosfat InsP_6) naturally found in wheat grain as soluble sodium and potassium salts (Torre *et al.*, 1991). During food processing of cereal products or during digestive stage InsP_6 is capable to fix metal cations and can form stable structures which are not able to diffuse through gastrointestinal wall (Official Methods of Analysis, 1990). One mmol of InsP_6 is able to fix up to 6 mmol of bivalent metallic cations (Fe, Ca, Zn) (Harland *et al.*, 1987). This contributes to an essential demineralization of food in the case of high intake of fibres and cereals.

During food fortification, processing and storage numerous physico-chemical and enzymatic processes take place which may greatly influence biological value of these products. Hereby, elaboration of a technologic process of fabrication of fortified products should be based on a profound study of the evolution of these micronutrients during technological process as well as during consumption.

The aim of the present study is investigation of the influence of bread making method on iron bioavailability from bread fortified with iron. The process of enzymatic degradation of phytates (InsP_6) during gastro-intestinal digestion and *in vitro* was studied as well.

The study of iron bioavailability was performed *in vitro*, according to the Monsen model and *in vivo* on white laboratory rats.

2. Materials and methods

During the study, we used bread made of superior quality wheat flour of autochthon production. As iron additive, iron sulphate (II) ($\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$) was used, which is accepted as food additive by “Codex Alimentarius” in the Republic of Moldova.

2.1. Bread making procedures

Bread was made by direct (mono-phase) method, which comprises the concurrent mixing of all the ingredients; indirect (bi-phase) method with the administration of additive at the kneading stage and by lactic-acid method.

To reproduce the fabrication of integral wheat bread, the integral reconstituted wheat flour was used (flour plus bran). For the preparation of bread with yeast 6 kg of integral wheat flour (4.65 kg flour plus 1.35 kg bran) were mixed with 3.6 l distilled water and 150 g of yeast *Saccharomyces cerevisiae*. For lactic-acid fermentation, the dough was prepared by the mixing of 1.2 kg integral wheat flour (0.93 kg flour plus 0.27 kg bran) with 600 ml distilled water. After one day of natural fermentation at 30°C, the dough was mixed with 4.8 kg of integral flour and 3 l of distilled water (Lopez *et al.*, 2003).

For bread prepared by lactic-acid method with yeast addition, the ingredients were mixed during 5 minutes. The obtained dough was stored at 30°C to allow fermentation to start. Then the dough was knitted again for the gluten formation and left at 30°C. The dough was backed after 3 hours of fermentation for the bread with yeast and one hour for dough made by lactic-acid method. Bread was lyophilized, milled and introduced in the alimentation of rats.

2.2. In vitro analysis procedures

Iron *in vitro* digestibility, as well as investigation of enzymatic degradation process of phytates was performed in two stages: gastric stage (pH=2, in pepsin presence) and intestinal stage (pH 8.2, in trypsin presence) (Moisen, 1991; Miller *et al.*, 1981).

A quantity of 10 g sample were stored at $37 \pm 0.1^\circ\text{C}$ for 15 min in acid environment (pH = 2) created by addition of 1.5 N HCl solution. After pepsin administration (150 mg/100g product) the blend was incubated for two hours at $37 \pm 0.1^\circ\text{C}$ with continuous mixing. Each 30 min equal samples were taken for the analysis, centrifuged for 10 min (6000 rot/min). The intestinal stage was realized in the same conditions after the establishment of the pH value at 8.2, in trypsin presence in NaHCO_3 0.08 M environments. The amount of soluble iron was determined through the spectrophotometric method (Kosse *et al.* 1996).

Phytic acid amount (InsP_6) in the flour and the product was determined through the spectrophotometric method (Official Methods of Analysis, 1990). The method is based on the organic phosphates capacity to interact with the ammonium molybdate and vanadate with formation of a yellowish – gold complex. The intensity of the colour is correlated with phosphorus amount. Subsequently the amount of InsP_6 was recalculated since phytic phosphor constitutes just 28.2% of the total InsP_6 .

The analysis were made using DR-5000 spectrophotometer, all tests were made in triplicates.

2.3. *In vivo analysis procedures*

Animals and their alimentation. To analyze the influence of the diet with different iron amounts (bread prepared by different procedures with iron addition) on the main haematological indexes the study on a sample of 20 white Wistar rats with the body weight of 180–210 g divided in 3 groups (6 – 7 in each group) was designed. The animals from the first group (control sample) were fed with ordinary non fortified bread (bread prepared by classical and bi-phase method). The second group comprised the rats fed with bread prepared through traditional, bi-phase method enriched with iron (bread prepared through the traditional, bi-phase method with 80 mg/kg iron addition). The third group comprised the rats fed with bread prepared by lactic-acid fermentation method enriched with iron (80 mg Fe/kg). Animals from all three groups were kept on the above mentioned diet for 21 days.

2.4. *Models of samples obtaining and analytic procedures*

Biological material – peripheral blood from the laboratory animals was obtained twice – initially, before the beginning of the experiment and at the end of the experiment through the procedure described by Ciudin *et al.* (1997).

Blood was collected to single use test tubes Ependorf type of 1.5 ml. Blood indexes have been appreciated in the heparinized peripheral blood in the haematological analyzer PCE-210 (ERMA, Japan).

Blood serum was obtained through centrifugation of the peripheral blood at 3000-4000 rot/min for 15 minutes. After centrifugation, blood serum was transferred in Ependorf test tubes and was stored in

the refrigerator at +4°C till the end of the experiment.

Blood iron level determination was performed using the method with cromasorol according to the working instructions of the analyzer (Iron Cromazurol, Eliteh, France).

3. Results and discussion

Production processes of cereal products (flour moistening, fermentation and dough knitting, backing and drying) influence greatly on the amount of phytates in food products. Hereby, vegetal phytase, an enzyme that degrades phytic acid and its soluble sodium and potassium salts, reduces essentially the amount of this antinutrient during technological processing of the flour especially during fermentation. This process is explained by the establishment of optimal conditions for development of phytase activity during dough fermentation at 35...40°C, pH value 4–5, as well as due to the fact that bread yeast amplify this process through their own phytase activity (Turk *et al.*, 1996).

However, recent use of fast bread making procedures, as well as the use of the iron supplements at the beginning of the dough fermentation leads to rapid fixation of bivalent metal cations in chelate compounds such as $\text{InsP}_6 \cdot x \text{Fe}$, subsequently not capable to diffuse the gastrointestinal wall and they are not subjected to enzymolysis.

In order to demonstrate the influence of bread making procedure (type and duration of fermentation, bread yeast quality), the results obtained during application of traditional bread making method – mono-phase and bi-phase method as well as of lactic-acid fermentation method based on the use of the wheat bran reconstituted through the addition of bran were compared (Table 1).

3.1. *In vitro procedures*

It was established that during mono-phase method, only 15% phytates presented in the bread were subjected to hydrolysis during 2 hours of *in vitro* gastro-intestinal digestion. Soluble iron amount (dialyzable) was, at the end of these 2 hours of gastro-intestinal digestion, 0.02 mg/100g product, which constitutes just 4.6% of the total iron naturally present in the product (2.5 mg/100g product).

Table 1. The influence of bread making procedure and the recipe on the iron bioavailability* in bread products**

Bread making method	Bread yeast	Fe dialyzable mg %			Fe dialyz. %	Soluble phytates, mg %			Soluble phytate %
		gastric	intestinal	Total		gastric	intestinal	Total	
Tradit. (mono-phase)	100%	0.07±0.02	0.05±0.01	0.12±0.02	4.6±0.5	51.3±1.9	33.2±1.6	84.5±1.9	14.6±0.4
Tradit. (bi-phase)	100%	0.15±0.01	0.06±0.01	0.21±0.01	8.1±0.4	125.2±1.7	89.5±1.3	214.7±1.7	36.5±0.3
Lactic-acid	–	0.29±0.02	0.25±0.02	0.54±0.02	21.6±0.8	217.5±2.2	116.8±1.9	334.3±2.2	56.8±0.4
	25%	0.32±0.03	0.24±0.02	0.56±0.03	22.4±1.2	212.6±2.7	87.5±1.6	300.1±2.7	51.0±0.5
	50%	0.26±0.02	0.08±0.01	0.34±0.02	13.6±0.8	197.9±2.5	99.7±2.2	297.6±2.5	50.6±0.4
	75%	0.22±0.01	0.24±0.02	0.46±0.02	18.4±0.8	195.5±2.9	131.3±3.1	326.8±3.1	55.6±0.5

III - Fe total – 2.5 ± 0.2 mg %

** phytates –total phytates amount in the product 578.0 ± 5.0 mg phytic acid / 100 g product

Phytates in the bread made by bi-phase method underwent hydrolysis in 36.5%. In this case, the amount of the dialyzable iron constituted 0.2 mg/100g product, or approximately 8% from the total iron amount.

Obviously, this difference is due to bread making procedure, because the ingredients composition is the same in all the cases. In the first case, fermentation procedure goes rapidly while during bi-phase procedure in the starter an important phytase activity will develop due to the phytase present in the flour as well as due to the enzymes present in the bread yeast. The administration of the flour at the dough formation stage is followed by a considerable phytate enzymatic hydrolysis, which leads to a more intense release of the iron from the insoluble phytates complexes.

A number of previous studies showed that lactic-acid fermentation of the dough, established using bran from the reconstituted wheat flour, contributes greatly to an essential decrease of the phytates amount (Lopez *et al.*, 2003). Lactic-acid microflora, isolated from the dough obtained through natural fermentation is capable to degrade phytates. Together, phytic degradation and formation of lactic acid could contribute to a more intense release of the minerals in the products.

Hereby, during this study the influence of lactic-acid fermentation, established on the fasts of bran from the reconstituted wheat flour, was investigated. In order to show the important role of yeast on this process, bread products were made. After lactic-acid fermentation development (with the duration of fermentation of 24 hours) into the recipe (prescription) for traditional (bi-phase) method respectively 25, 50 and 75% of yeast was added.

According to obtained results, in the case of application of the natural dough fermentation procedure, 50-57% of the present phytates were

hydrolyzed after 2 hours *in vitro* gastro-intestinal digestion. The addition of the bread yeast does not seem to have an important effect (table 1). This shows that even though phytate decrease during traditional bread making procedure was identified, this cannot be referred to the phytase activity of the bread yeast but is due to endogen flour phytase activity.

Dialyzable iron amount in this case is higher (0.34–0.56 mg/100g product), which constitutes 13–22% from the total iron present in the product. This is obviously due to an important degradation of phytates, which makes this mineral insoluble. Apparently, the naturally present iron intake from wheat flour is insufficient for diets, which comprise few products that contain haemic iron. The study of iron bioavailability in the fortified bread products, prepared through the bi-phase and lactic-acid method shows that in the case of bi-phase method only 8–12 % of the present iron is presented in its soluble form after 2 hours of gastro-intestinal digestion (Table 1). In the case of bread prepared by lactic-acid method, the rate of the dialyzable iron constitutes 20–22 %.

It is obvious that *in vitro* investigations do not reflect the multitude of the factors that may influence the mineral valorisations in the gastro-intestinal tract. First, there are subjective factors that reflect body reserve of iron. According to the literature data, *in vitro* bioavailability is related with *in vivo* digestion in a proportion of 60–76 % (Lopez *et al.*, 2003, Miller *et al.*, 1981).

3.2. *In vivo* procedures

Serum iron concentration is an indicator of the amount of iron in plasma, fixed to a specific protein – transferrin. The majority of iron which is part of plasma derives through catabolism of reticulo-endotelial system, and iron that leaves plasma derives from bone marrow. Regardless the

fact that plasmatic iron reserve is not significant (approximately 3 mg), the last is very active because it transports daily about 30 mg of iron. Transport parameters of the generally are not modified till body reserve of iron is not completely depleted. In the convalescence period, after a severe anaemia, namely serum iron is indicator of recovery degree and of the body iron statute.

It was established, in the experimental groups, that after the administration of the special diet the level of serum iron was of $36.48 \pm 13.8 \mu\text{mol/L}$ for the control group (I), of $41.55 \pm 19.58 \mu\text{mol/L}$ (II group) and of $40.14 \pm 18.78 \mu\text{mol/L}$ for group III (Figure 1).




Figure 1. Serum iron evolution after the application of the special food: I-control group; II- fortified bread produced through traditional method; III- fortified bread produced by lactic-acid method

The investigations reveal that iron intake and procedures applied for bread making have a significant importance for bioavailability of this essential mineral. Food products rich in fibers such as products made of integral wheat flour contain an important amount of phytates, which may reduce essentially iron absorption and utilization ability in the human body. Antinutritive effect of the phytic acid from cereals is manifested through its ability to chelate metal ions (Fe^{3+} , Zn^{2+} , Mg^{2+} , Ca^{2+}), at gastro-intestinal pH value thus forming insoluble complexes metal-phytate that makes the metal not available for absorption in the gastro-intestinal tract (Carpenter *et al.*, 1992; Guthrie *et al.*, 1995). Phytic acid also interacts with some enzymes (trypsin, pepsin, α -amylase, β -galactosidase), proteins and vitamins reducing their activity. It was established that phytates suppress activity of α -amylase through the chelating of calcium necessary for the enzymatic activity. With proteins, phytic acid forms complexes protein-phytates or protein-mineral-phytates which leads to a decrease of the solubility, digestibility and functionality of the proteins (Mothes *et al.*, 1988). Dephytinisation (phytate extraction), performed through different

procedures, increases significantly iron absorption (Mothes *et al.*, 1988; Hallberg, 1995a).

The procedures applied for the food transformation (knitting, moistening, fermentation, and backing) have an important influence on the amount of phytic acid in the final product, which may be the subject of enzymatic hydrolysis due to endogen phytase presence as well as due to exogen phytase from yeast. An essential role play duration and food transformation conditions (Hallberg, 1995b; Brune *et al.*, 1992; Hallberg, 1980). Phytase activity is optimal at pH value 5–5.5 and temperature 55°C , the enzyme being still active at 70°C (Tseng M., *et al.*, 1997). In the dough, together with endogen phytase of the dough, the phytase from yeast may also influence if it is added (Banu, 2000). Phytic acid and phytate hydrolysis takes place at all stages of the bread making process, being influenced by: flour extraction degree, the amount of yeast used; temperature and duration of fermentation; dough pH value; water amount in the dough, phytate solubility especially of its salts; additive presence such as ascorbic acid and sodium bicarbonate (Banu, 2000). More than that, phytase has a significant role for the phytate hydrolysis in the stomach and intestine, thus possible decrease of mineral absorption due to phytic acid resistance for digestion in are human intestine should be taken into consideration.

Numerous studies have shown that lactic-acid bacteria are able to degrade phytic acid, and lactic-acid fermentation increases Ca, Mg, Fe solubility as estimated during *in vitro* investigations (Lopez *et al.*, 2003, Shirai *et al.*, 1994). All together, phytic acid degradation and the lactic acid formation have improved mineral bioavailability in the bread prepared lactic-acid method.

In this study, it was established that non-hem iron from the bread prepared by lactic-acid fermentation method has an increased *in vitro* bioavailability compared to the bread prepared by traditional methods (mon- and bi-phase). In the same time, a decrease of the amount of phytic acid during *in vitro* gastro-intestinal digestion was noticed. These observations show that there is a direct relationship between potential iron bioavailability from the bread products and phytase degradation degree.

During *in vivo* investigations it was established that for the rats from all 3 groups the parameters of peripheral blood after 21 days of special diet administration were within established limits. For the II and III group, where the special diet was

used (fortified bread prepared by traditional method and fortified bread prepared by lactic-acid method) an essential increase of haemoglobin, erythrocyte number, platelets number was observed compared to the control group. This is obviously due to an increased iron intake from the administered special diet. But the most significant influence was noticed in the case of the supplement administration on the serum iron level, which reflects body iron reserve. For those 2 experimental groups, fed with fortified bread, body iron reserve is considerably higher compared to the control group.

The comparison of biochemical indices of the blood for those two rat groups fed with fortified bread did not allow us to reveal the influence of bread making method. Hereby, although *in vitro* investigations showed that iron dialysability from the bread prepared by lactic-acid method is considerably higher ($\approx 20\%$) compared to the iron dialysability from traditionally prepared bread ($\approx 8\%$), *in vivo* investigations did not testify any significant difference for the biochemical indices for those two experimental groups.

The reason might probably be due to the considerable iron intake, which allowed to satisfy the requirements and to complete the body iron reserve. An essential role plays by the initial iron statute of those 3 rat groups that was within the normal limits at the beginning of the experiment.

4. Conclusions

Our investigations established that procedure applied for fortified bread production has a key influence on iron bioavailability degree in *in vitro* gastro-intestinal conditions. This fact is directly related with an advanced level of the phytate degradation from the bread, due to applied bread making procedure and also due to a longer bread-making procedure.

The study of biochemical blood indices, in samples collected from the laboratory animals fed with fortified bread (8 mg/100g product), that was prepared by traditional and by lactic-acid methods compared to the control group, showed that the key role in animal nutrition plays iron intake. Hereby, iron statute of both experimental groups especially body iron reserve was significantly improved compared to the control group.

In case of sufficient iron consumption bread making procedure did not have any influence on

iron statute in examined groups. In order to show the influence of bread-making procedure on iron status, it is necessary to investigate the influence of special diet administration on the laboratory animals with induced anaemia.

On the basis of the received experimental results it has been elaborated an algorithm, allowing to determine the iron absorption coefficient from a representative meal and the estimation of the providing degree of nutritional requirements for different groups of population.

References

- Banu C. *et al.* 2000. *Biotehnologii în industria alimentară (Biotechnologies in food industry)* Editura Tehnică, București.
- Bauernfeind J.C. 1988. Nutrification of food. In *Modern nutrition in Health & Disease*, 7th Ed.M.E. Shils and Y.R. Young (eds). Lea & Febiger, Philadelphia.
- Brune M., Rossander-Hulten L., Hallberg L., Gleerup A. & Sandberg A.S. 1992. Iron absorption from bread in humans: inhibiting effects of cereal fiber, phytate and inositol phosphates with different numbers of phosphate groups. *J. Nutr.* **122**: 442–449.
- Carpenter C.E. & Mahoney A.W. 1992. Contributions of Heme and Nonheme Iron to Human Nutrition. *Crit. Rev. Food. Sci. Nutr.* **31**: 333–367.
- Ciudin E. & Marinescu D. 1997. *Patologia animalelor de laborator și tehnică experimentală*. Editura Moldogrup, Iași, p. 203 – 205.
- Guthrie H.A. & Picciano M.F. 1995. *Human nutrition*. Ville Saint-Louis, Mosby.
- Hallberg L. 1980. Food iron absorption. In : Cook JD, ed. *Methods in hematology*. Vol I. London: Churchill, 116–133.
- Hallberg L. 1995a. Results of surveys to assess iron status in Europe. *Nutr. Rev.* **53**: 314–322.
- Hallberg L. 1995b. Iron balance in menstruating women. *Eur. J. Clin. Nutr.* **49**: 200–207.
- Harland B.F. & Oberleas D. 1987. Phytate in foods. *Wld. Rev. Nutr. Diet.*, pp. 52, 235–259.
- Kosse J.S., Young A.C., Gil A.I. & Miller D.D. 2002. A rapid method for iron determination in fortified foods. *Food Chemistry* **75**: 371–376.
- Lopez H.W., Duclos V., Coudray Ch. *et al.* 2003. Making bread with sourdough improves mineral bioavailability from reconstituted whole wheat flour in rats. *Nutrition* **19**: 524–530.
- Luten J., Crews H. & Flynn A. 1996. Interlaboratory trail on the determination of the *in vitro* iron

- dialysability from food. *J. Sci. Food Agric.*, **72**: 415–424.
- Miller D.D., Schrickler B.R. & Rasmussen R.R. 1981. An in vitro estimation of iron availability from meals. *Am. J. Clin. Nutr.*, **34**: 2248–2256.
- Moisen S. 1991. A model for feed evaluation based on in vitro digestible dry matter and protein. *CAB International*, Vellingford, UK, pp. 135–145.
- Mothes R., Schwenke K.D., Lizwer Z.D., Gast K. & Welfle H. 1988. Studies on rapeseed protein – phytic acid interaction. *Acta alim.*, 17.4 pp. 364–365.
- Motruc N., Ciobanu C., Deseatnicov O. & Sturza R. 2005. Aportul nutrițional în micronutrimenți în alimentația copiilor instituționalizați și posibilități de optimizare. *Proceedings of the 30th Annual Congress of the American Romanian Academy of Arts and Sciences (ARA)*, July 5–10, 2005.
- Shirai K., Revah-Moiseev S., Garcia-Garibay M. & Marshall V.M. 1994. Ability of some strains of lactic acid bacteria to degrade phytic acid. *Lett. Appl. Microbiol.*, **19**: 366.
- Torre M. & Rodriguez A.R. 1991. Effects of dietary fiber and phytic acid on mineral availability. *Crit. Rev. Food Sci. Nutr.*, 1, pp. 1–22.
- Tseng M., Chakraborty H., Robinson D.T., Mendez M, & Kohlmeyer L. 1997. Adjustment for iron intake for dietary enhancers and inhibitors in population studies: bioavailable iron in rural and urban residing Russia women and children. *J. Nutr.*, **127**: 1456–1468.
- Turk M., Carlsson N.G. & Sandberg A.S. 1996. Reduction in the levels of phytate during wholemeal bread making: effect of yeast and wheat phytases. *J. Cereal Chi.*, **23**: 257.
- * * * Official Methods of Analysis, fifteenth ed., 1990, Vol. II, p. 800.
- * * * Raportul UNICEF “Alimentația și nutriția umană în Republica Moldova”, Biroul pentru Moldova, 2002.
- * * * Starea de nutriție în Republica Moldova. Raport UNICEF 1996–2000, p. 38.