



Two-Dimensional Superconducting Fluctuations in MgB₂ Films

Sidorenko A. S., Zdravkov V. I., Ryazanov V. V., Klemm M., Horn S., Tidecks R., Müller C., Wixforth A.

<https://doi.org/10.1023/B:JOSC.0000021215.50054.7d>

Abstract

The origin of the resistive transition broadening for superconducting MgB₂ films is investigated experimentally. The crucial role of two-dimensional weak and critical fluctuations is demonstrated.

References

1. J. Namagatsu, N. Nagakawa, T. Muranaka, Y. Zenitani, and J. Akimitsu, Nature 410, 63(2001).
2. V. V. Schmidt, Introduction to the Physics of Superconductors, 2nd Russian edn., enlarged by V. V. Ryazanov and M. F. Feigelman (MCNMO Publishers, Moskow, 2000) Chapter III, § 19.
3. N. Ya. Fogel, A. S. Sidorenko, L. F. Rybalchenko, and I. M. Dmitrenko, Sov. Phys. JETP 50, 120(1979).
 - A. Kapitulnik, Physica C 153-155, 520(1988).
 - A. S. Sidorenko, L. R. Tagirov, A. N. Rossolenko, N. S. Sidorov, V. I. Zdravkov, V. V. Ryazanov, M. Klemm, S. Horn, and R. Tidecks, JETP Lett. 76, 17(2002).
4. L. G. Aslamazov and A. I. Larkin, Phys. Lett. 26A, 238(1968).
5. L. P. Kadanoff and G. Lamore, Phys. Rev. 175, 579(1968).