

CMT 4P THE INFLUENCE OF THE RASHBA SPIN-ORBIT COUPLING ON TWO-DIMENSIONAL MAGNETOEXCITON-POLARITONS IN MICROCAVITY

E. V. Dumanov¹, S. A. Moskalenko¹, I. V. Podlesny¹, M. A. Liberman^{2,3}, S. S. Rusu¹,
V. M. Bajireanu¹

¹*Institute of Applied Physics of the Academy of Sciences of Moldova, Academic Str. 5,
Chisinau2028, Republic of Moldova*

²*Nordita, KTH Royal Institute of Technology and Stockholm University, Roslagstullsbacken23,
10691 Stockholm, Sweden*

³*Moscow Institute of Physics and Technology, Institutskii per. 9, Dolgoprudnyi,
Moskovsk. obl.,141700, Russia*

Our results concern the electron-radiation and Coulomb electron-electron interactions in system of two-dimensional (2D) electron-hole pairs on the layer embedded into the microcavity and subjected to the action of a strong perpendicular magnetic field as well as of the electric field in the same direction giving rise to the Rashba spin-orbit coupling (RSOC) with the third order chirality terms and with a quartic term in the nonparabolic dispersion law of the heavy holes in the GaAs-type quantum well.

In these conditions the magnetoexciton-polariton branches have the nonmonotonous dependences on the magnetic field strength B with intersections and strong changing of the quantum transition probabilities. The more simple cases without polariton branches were described earlier in the Refs. [1, 2].

The Rabi frequencies which determine the spread of the polariton branches are proportional to the dipole moment P_{cv} of the optical quantum transition between the valence and conduction bands, they increase as \sqrt{B} with the increasing of the magnetic field and obey to the selection rule following which the numbers of the Landau quantization levels of the electron and hole in the composition of the magnetoexciton coincide. Similar properties were revealed in Ref. [3].

The coherent macroscopic state was introduced into the Hamiltonian following the Keldysh-Kozlov-Kopaev method in combination with the Bogoliubov theory of the quasiaverages. The equations of motion were deduced for the two-particle integral operators $\rho(\vec{Q})$ and $D(\vec{Q})$ describing the optical and acoustical plasmons correspondingly as well as for the creation and annihilation operators $\hat{\psi}_{ex}^\dagger(\vec{k}_\parallel)$, $\hat{\psi}_{ex}(\vec{k}_\parallel)$, $C_{\vec{k}_\tau, \vec{k}_\parallel}^\dagger$, $C_{\vec{k}_\tau, \vec{k}_\parallel}$ of the magnetoexciton and photon modes in the microcavity correspondingly. Such possibility is due to the degeneracy of the Landau levels, which in Landau gauge do not depend on the one-dimensional wave vector. It was shown that the presence of the SOC is equivalent to take into account the influence of some excited Landau levels. It permits to obtain following [4] the energy spectrum of the collective elementary excitations, even when the ground state of the system is the Bose-Einstein condensation of the 2D magnetoexciton-polaritons on the lower polariton branch with the in-plane vector $\vec{k}_\parallel = 0$. The calculations in this direction are continued.

E. V. D., I.V.P., and V.M.B. thank the Foundation for Young Scientists of the Academy of Sciences of Moldova for financial support (14.819.02.18F).

[1] S.A. Moskalenko et al, *Solid State Communications*, 151, (2011) , 1690–1695

[2] I.V. Podlesny et al, *Physica E*, 49, (2013), 45–51

[3] S.A. Moskalenko et al, *J. Nanophoton.*, 6(1), (2012), 061806

[4] S.A. Moskalenko et al, *Eur. Phys. J. B*, 85: 359 (2012)